

POSTER PRESENTATION

Open Access


Modeling Na^+ - and Ca^{2+} -dependent mechanisms of rhythmic bursting in excitatory neural networks

Ilya A Rybak^{1*}, Patrick E Jasinski¹, Yaroslav I Molkov^{1,2}, Natalia A Shevtsova¹, Jeffrey C Smith³

From Twenty First Annual Computational Neuroscience Meeting: CNS*2012
Decatur, GA, USA. 21-26 July 2012

The mechanisms generating neural oscillations in the mammalian brainstem, particularly in the pre-Bötzinger complex (pre-BötC) involved in control of respiration, and the spinal cord (e.g. circuits controlling locomotion) that persist after blockade of synaptic inhibition, remain poorly understood. Experimental studies in medullary slices from neonatal rodents containing the pre-BötC identified two mechanisms that could potentially contribute to generation of rhythmic bursting in the pre-BötC: one based on the persistent sodium current (I_{NaP}) [1,2], and the other involving the voltage-gated calcium (I_{Ca}) [3] and/or the calcium-activated nonspecific cation current (I_{CAN}), activated by intracellular Ca^{2+} accumulated from extra- and/or intracellular sources [4]. However, the involvement and relative roles of these mechanisms in rhythmic bursting are still under debate.

In this theoretical/modeling study we investigated Na^+ - and Ca^{2+} -dependent bursting generated in single cells and in a heterogeneous population of synaptically interconnected excitatory neurons with I_{NaP} , and I_{Ca} randomly distributed within the population. We analyzed the possible roles of network connections, ionotropic and metabotropic synaptic mechanisms, intracellular Ca^{2+} release, and the Na^+/K^+ pump in rhythmic bursting activity generated under different conditions. We show that the heterogeneous population of excitatory neurons can operate in different oscillatory regimes with bursting dependent on I_{NaP} and/or I_{CAN} , or independent of both (Fig. 1). The oscillatory regime and operating bursting mechanism may depend on neuronal excitability,



synaptic interactions and relative expression of particular ionic currents.

The existence of multiple oscillatory regimes and their state-dependency may provide explanations for different rhythmic activities observed in the brainstem and spinal cord under different experimental conditions.

* Correspondence: rybak@drexel.edu

¹Department of Neurobiology and Anatomy, Drexel University College of Medicine, Philadelphia, PA 19129, USA

Full list of author information is available at the end of the article

Author details

¹Department of Neurobiology and Anatomy, Drexel University College of Medicine, Philadelphia, PA 19129, USA. ²Department of Mathematical Sciences, Indiana University – Purdue University Indianapolis, IN 46202, USA. ³Cellular and Systems Neurobiology Section, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD 20892, USA.

Published: 16 July 2012

References

1. Butera RJ, Rinzel JR, Smith JC: **Models of respiratory rhythm generation in the pre-Bötzinger complex: II. Populations of coupled pacemaker neurons.** *J Neurophysiol* 1999, **82**:398-415.
2. Koizumi H, Smith JC: **Persistent Na⁺ and K⁺-dominated leak currents contribute to respiratory rhythm generation in the pre-botzinger complex in vitro.** *J Neurosci* 2008, **28**:1773-1785.
3. Pace RW, Mackay DD, Feldman JL, Del Negro CA: **Inspiratory bursts in the preBötzinger complex depend on a calcium-activated non-specific cation current linked to glutamate receptors in neonatal mice.** *J Physiol* 2007, **582**(Pt 1):113-125.
4. Thoby-Brisson M, Ramirez JM: **Identification of two types of inspiratory pacemaker neurons in the isolated respiratory neural network of mice.** *J Neurophysiol* 2001, **86**:104-112.

doi:10.1186/1471-2202-13-S1-P38

Cite this article as: Rybak *et al.*: Modeling Na⁺- and Ca²⁺-dependent mechanisms of rhythmic bursting in excitatory neural networks. *BMC Neuroscience* 2012 **13**(Suppl 1):P38.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

